



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C11D 1/825, 1/94, 1/835, A61K 7/50	A1	(11) International Publication Number: WO 99/19438 (43) International Publication Date: 22 April 1999 (22.04.99)
(21) International Application Number: PCT/US98/21788 (22) International Filing Date: 15 October 1998 (15.10.98) (30) Priority Data: 08/943,675 15 October 1997 (15.10.97) US (63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Application US 08/943,675 (CIP) Filed on 15 October 1997 (15.10.97) (71) Applicant (for all designated States except US): STEPAN COMPANY [US/US]; 22 West Frontage Road, Northfield, IL 60093 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): KNOX, Steven, J. [US/US]; Apartment #502, 555 West Arlington Place, Chicago, IL 60614 (US). MALIK, Arshad [US/US]; 3172 Riviera Way, San Ramon, CA 94583 (US). (74) Agent: NOONAN, Kevin, E.; McDonnell Boehnen Hulbert & Berghoff, Suite 3200, 300 South Wacker Drive, Chicago, IL 60606 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: HIGH FOAMING DETERGENT COMPOSITION HAVING NON-IONIC SURFACTANT BASE		
(57) Abstract <p>Disclosed are aqueous liquid cleaning compositions, the compositions being free of anionic surfactants and comprising: (a) linear alcohol ethoxylate; (b) amine oxide or betaine; and other, optional components, such as a cationic ammonium compound.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

HIGH FOAMING DETERGENT COMPOSITION HAVING NON-IONIC SURFACTANT BASE

Background of the Invention

5 This application is a continuation-in-part of International Application PCT/US97/06211 filed April 14, 1997, which is a continuation-in-part of U.S. Patent Application Serial No. 08/631,938, filed April 15, 1996.

10 Field of the Invention

 This invention relates to formulations for manually washing dishes, hand soaps, and for high foaming cleaning compositions.

Description of the Related Art

15 Light-duty liquid detergent formulations for kitchen surfaces are well known. Kitchen surfaces include counter tops, stove tops, dishes and any other hard surface commonly found in kitchen environments. The term "dishes" includes any utensils involved in food preparation or consumption. Kitchen surfaces,
20 particularly dishes, must be washed free of food residues, greases, proteins, starches, gums, dyes, oils and burnt organic residues.

 Most of the consumer accepted formulations in use include anionic synthetic surfactants with or without a nonionic
25 surfactant. Many of such formulations contain a sulfonate-type anionic surfactant, for example, an alkylbenzene sulfonate or an alkane sulfonate, in conjunction with a sulfate or alkyl ether sulfate, or a nonionic surfactant, for example, an alcohol ethoxylate, an alkyl phenol ethoxylate, a mono- or
30 diethanolamide or an amine oxide. The sulfonate material generally predominates.

 It is the anionic surfactant that provides the typical high foaming (suds) characteristics generally associated with dish washing formulations. Foam (suds) is the cleaning
35 efficacy signal relied on by consumers. Nonionic surfactants

generally do not provide good foaming characteristics.

U.S. Patent No. 2,746,928 discloses that it is not possible to mix anionic surface-active agents with quaternary ammonium germicides. The cationic quaternary ammonium
5 germicide reacts with the anionic surface-active agent resulting in a reduction in germicidal and detergent activity.

Thus, anionic surfactants are incompatible with cationic quaternary antimicrobial surfactants and nonionic surfactants do not normally provide significant foaming capability to
10 liquid formulations. Therefore, dish washing liquids combining good foaming and antimicrobial activities are not available to the consumer.

Solutions to the problems posed by the incompatibility of cationic and anionic surfactants have focused on various non-
15 ionic surfactant systems. While having the potential to overcome the known compatibility problems, such systems are not capable of the cleaning efficacy and foam volume demanded by consumers. Hence, there remains a critical need for cleaning compositions based on non-ionic surfactant systems that provide
20 excellent cleaning with high foam volume.

SUMMARY OF THE INVENTION

In general, anionic surfactant systems such as those found in the current light duty liquids are classified as high foamers. Conversely, nonionic surfactant systems are
5 classified as low foamers.

By careful selection and extensive experimentation, we have identified nonionic and nonionic/amphoteric surfactant mixtures that produce consumer acceptable foam comparable to commercial dish washing liquids that use anionic detergents.
10 The useful nonionic surfactants include ethoxylates that have various chain lengths not exceeding 12 carbon atoms and degrees of ethoxylation that allow the dish washing liquid to be effective on a wide range of food soils while providing excellent flash foam volume and foam stability. This system
15 provides the consumer with effective cleaning on, but not limited to, greasy food soils, fatty food soils, and oily food soils.

The invention provides surfactant compositions based on nonionic surfactant components that function as cleaning
20 compositions. Further included in the invention are disinfectant hand soaps, body washes, disinfectant or antibacterial dishwashing liquids, and conditioning shampoos. Each of these latter cleaning compositions includes at least one cationic ammonium salt. The specific cationic salts are
25 selected depending on the ultimate use or function of the cleaning composition.

Certain formulations of this invention will control the presence and spread of bacteria on hard surfaces in the kitchen environment, especially dishes. In this context, the invention
30 is a microbiological active quaternary ammonium salt ingredient homogeneously incorporated into a nonionic aqueous surfactant system. Unexpectedly, the formulations of the invention have excellent flash foaming and residual foaming capability although no anionic surfactants are included.

The invention also provides personal care compositions including a quaternary ammonium compound which is a conditioning compound.

Thus, the invention provides hand soap compositions comprising a nonionic surfactant base in combination with at least one cationic ammonium compound. The ammonium compound may be an antibacterial compound or a conditioning agent, or both. Certain hand soap formulations will include a conditioning agent and an antimicrobial compound. Similar formulations may be formulated to function as conditioning shampoos.

The unexpected foaming properties of the formulations of the invention are illustrated in the examples. The foaming properties are due to the carefully balanced mix of nonionic surfactants. The formulations tested in these examples contain preferred concentrations of ingredients.

Thus, the invention provides aqueous liquid cleaning compositions, the compositions being free of anionic surfactants and consisting essentially of a nonionic surfactant system and a cationic ammonium compound.

Significantly, the invention also provides high foaming nonionic or nonionic/amphoteric systems that are excellent grease cutters.

The nonionic surfactant system may comprise (1) from about 0.1-50% by weight based on the weight of the composition of a linear alcohol ethoxylate having an average carbon chain length of no more than 12 carbon atoms; and (2) a surfactant member selected from the group consisting of amine oxides and betaines. In these compositions, the total concentration of active components in the composition based on the weight of the composition is at least about 5%. Optional non-ionic surfactants include alkanolamides, alkyl polysaccharides, betaines, and polyhydroxy fatty acid amides. In various embodiments of the invention, these optional components may

replace a portion of the alcohol ethoxylate.

The nonionic surfactant systems of the invention may be combined with a variety of cationic ammonium compounds, such as for example, quaternary ammonium compounds or cationic
5 conditioning agents, to produce a cleaning composition that functions as dishwashing cleaner such as a an antimicrobial dishwashing liquid or handsoap or as a conditioning cleaner such as a conditioning shampoo.

Formulations of the invention can be utilized as body
10 washes and soaps, facial cleanser, bath gels, and bubble bathes.

In addition to providing styling and conditioning to the hair, the inventive formulations provide for repair of split ends.

DETAILED DESCRIPTION OF THE INVENTION

As used herein, the term disinfecting or disinfectant refers to antimicrobial and/or antibacterial activity. Disinfectant, antimicrobial and antibacterial formulations of the invention are capable of reducing the rate of microbial, i.e., bacterial, reproduction, and/or killing microbial organisms.

The invention encompasses detergent compositions containing various combinations of linear alcohol ethoxylates and nonionic surfactants selected from amine oxides and betaines. Typical compositions also include at least one cationic ammonium compound. In preferred embodiments, the detergent or cleaning compositions comprise a linear alcohol ethoxylate, an amine oxide, an alkyl mono- or dialkanolamide, and a cationic ammonium compound. In such compositions, the balance of the material is water. In particularly preferred embodiments, the weight ratio of linear alcohol ethoxylate to amine oxide is from about 3:1 to 1:3.

A particularly preferred detergent composition according to the invention is the following:

a detergent formulation free from anionic surfactants consisting essentially of:

(a) from about 2-23%, preferably 8-18%, by weight of a linear alcohol ethoxylate having 6-12 carbon atoms and 3-12, preferably 3 to 7 moles of ethoxylation per mole of alcohol;

(b) from about 2-23%, more preferably 4-23%, by weight of an amine oxide selected from the group consisting of (C₈₋₁₆) alkyl amido (C₁₋₄) alkyl di(C₁₋₄) alkyl amine oxides and (C₁₀₋₁₆) alkyl amine oxides; and optionally

(c) from about 1-10%, more preferably 3-7%, by weight of (C₁₀₋₁₆) alkyl mono- or dialkanolamides, where each alkanol portion independently has from 1-6 carbon atoms; and

(d) from about 0.5 to 20% of a cationic ammonium compound. In such compositions, the weight ratio of

component (a) to component (b) is most preferably from about 1:3 to 3:1. Most preferred linear alcohol ethoxylates have about 4.5 moles of ethoxylation per mole of alcohol. Most preferred alkanolamides are present at about 4-6% by weight of
5 the composition.

Another particularly preferred composition of the invention is a liquid cleaning composition consisting essentially of, by weight of the composition, from about 4-7% of a C₈₋₁₀ alcohol ethoxylate having an average of about 9 moles
10 of ethylene oxide, from about 12-20% of a C₈₋₁₀ alcohol ethoxylate having an average of about 12 moles of ethylene oxide, from about 7-13% of a fatty acid amidopropylamine oxide having an average of about 10-18 fatty acid carbon atoms, from about 1-4% of a fatty acid diethanolamide having an average of
15 about 10-18 fatty acid carbon atoms, from about 1-4% of a fatty acid monoethanolamide having an average of about 10-18 fatty acid carbon atoms; and an antibacterial effective amount of an antibacterial quaternary ammonium compound. A preferred
20 antibacterial quaternary ammonium compound is an alkyl dimethyl benzyl ammonium chloride. The balance of the composition is water. Such a composition may also contain an emulsifier or thickener such as xanthan gum, as well as fragrances, etc.

Still another particularly preferred formulation according to the invention is an aqueous liquid cleaning composition, the
25 compositions being free of anionic surfactants and consisting essentially of:

(a) from about 13-19% by weight based on the weight of the composition of a linear alcohol ethoxylate having an average carbon chain length of no more than 12 carbon
30 atoms; and;

(b) from about 3-7% by weight of the composition of a mono- or dialkanolamide; and

(d) from about 5-10% by weight of the composition of an alkylpolyglycoside.

Yet another composition consists essentially of:

- (a) from about 13-19% by weight based on the weight of the composition of a sulfobetaine;
- (b) from about 5-20% by weight of the composition of an amine oxide, a betaine, or mixture thereof;
- (c) from about 3-7% by weight of the composition of a mono- or dialkanolamide; and
- (d) from about 3-7% by weight of the composition of an alkylpolyglycoside,

the total concentration of surfactants in the composition being from about 30-35% by weight of the composition.

Another preferred aspect of the invention is a detergent composition suitable for preparing an aqueous, liquid cleaning formulation, the detergent composition comprising, based on the total weight of active components in the composition:

- (a) from about 10-70% by weight of an alcohol ethoxylate having an average of 6-12 carbon atoms and an average of 4-15 moles of ethylene oxide;
- (b) from about 15-60% by weight of an amine oxide, a betaine or mixture thereof;
- (c) from about 5-20% by weight of an alkanolamide having an average of 10-16 carbon atoms.

In yet another preferred aspect, the invention provides a detergent composition suitable for preparing an aqueous, liquid cleaning formulation, the detergent composition comprising, based on the total weight of active components in the composition:

- a) from about 25-60% by weight an alcohol ethoxylate having an average of 6-12 carbon atoms and containing an average of from about 4-15 moles of ethylene oxide where the concentration of alcohol ethoxylate in the

composition is of the total surfactant present by weight;

- b) from about 15-55% by weight of an amine oxide, a betaine or mixture thereof; and
- 5 c) 5-20% by weight of an alkanolamide having an average of from 10-16 carbon atoms; and
- d) up to about 20% by weight of a cationic surfactant.

10 Another preferred aspect of the invention is an aqueous cleaning formulation comprising from about 5% to 60% by weight of the formulation of a detergent composition,

the detergent composition comprising, based on the total weight of active components in the detergent composition:

- 15 (a) from about 25-60% by weight of an alcohol ethoxylate having an average of 8-10 carbon atoms and containing an average of 5-12 moles of ethylene oxide;
- (b) from about 15-55% by weight of an alkyl dimethyl amine oxide, an alkyl amido propyl dimethyl amine oxide, an alkyl amido propyl betaine, or a mixture
20 thereof;
- (c) from about 9-16% by weight of an alkanolamide having an average of 10-16 carbon atoms; and
- (d) from about 3-10% by weight of biocidal cationic
25 surfactant.

Optional, non-essential ingredients include fragrances, dyes, stabilizers, thickeners, etc.

Nonionic surfactants

30 The surfactants suitable for use in the inventive compositions include the following nonionic surfactants.

Alcohol ethoxylates

In the condensation products of aliphatic alcohols with ethylene oxide, i.e., alcohol ethoxylates, the alkyl chain of

the aliphatic alcohol can either be straight or branched and generally contains from about 5 to about 22 carbon atoms. The chain of ethylene oxide can contain from 2 to 30 ethylene oxide moieties per molecule of surfactant. Examples of such ethoxylated alcohols include the condensation product of about 5 6 moles of ethylene oxide with 1 mole of tridecanol, myristyl alcohol condensed with about 10 moles of ethylene oxide per mole of myristyl alcohol, the condensation product of ethylene oxide with coconut fatty alcohol wherein the coconut alcohol is 10 a mixture of fatty alcohols with alkyl chains varying from 10 to 14 carbon atoms and wherein the condensate contains about 6 moles of ethylene oxide per mole of alcohol, and the condensation product of about 9 moles of ethylene oxide with the above-described coconut alcohol. Examples of commercially 15 available nonionic surfactants of this type include Tergitol 15-S-9 marketed by the Union Carbide Corporation, Neodol 23-7 marketed by the Shell Chemical Company and Kyro EOB marketed by the Procter & Gamble Company.


A preferred concentration for the alcohol ethoxylate in 20 the composition is from about 10-70% of the total active surfactant by weight; i.e., it is from about 10-70% by weight based on the total weight of active components. A more preferred range is 25-60% by weight. A most preferred range is 40-60% by weight.

25 Generally, the alcohol ethoxylates have an average of 6-12 carbon atoms and contain an average of about 4-15 moles of ethylene oxide. Preferably, the alcohol ethoxylates have an average of 8-10 carbon atoms and an average of about 5-12 moles ethylene oxide. The most preferred alcohol ethoxylates 30 are selected from C_{9-11} alkyl ethoxylate (8EO), C_{6-10} alkyl ethoxylate (4EO), C_{8-10} alkyl ethoxylate (9EO), C_{8-10} alkyl ethoxylate (12 EO), C_{11} alkyl ethoxylate (5 EO), and a C_{11} alkyl ethoxylate (7EO).

Amide Surfactant

The amide type of nonionic surface active agents includes the ammonia, monoethanol and diethanolamides of fatty acids having an acyl moiety of from about 7 to about 18 carbon atoms. These acyl moieties are normally derived from naturally occurring glycerides, e.g., coconut oil, palm oil, soybean oil and tallow, but can be derived synthetically, e.g., by the oxidation of petroleum, or by the Fischer-Tropsch process.

The amide surfactants useful herein may be selected from those aliphatic amides of the general formula:



wherein R^4 is hydrogen, alkyl, or alkylol and R^5 and R^6 are each hydrogen, C_2 - C_4 alkyl, C_2 - C_4 alkylol, or C_2 - C_4 alkylenes joined through an oxygen atom, the total number of carbon atoms in R^4 , R^5 and R^6 being from about 9 to about 25. A further description and detailed examples of these amide nonionic surfactants are contained in U.S. Pat. No. 4,070,309, issued to Jacobsen on Jan. 24, 1978. That patent is hereby incorporated herein by reference.


Preferred amide surfactants have an average of from 10-16 carbon atoms in the portion of the molecule contributed by the fatty acid. More preferably, the amide surfactants have an average of from 8-10 carbon atoms in the fatty portion. Most preferred amide surfactants are lauric (average of C_{12}) myristic (average of C_{14}) mono ethanolamide, lauric diethanolamide, lauric monoethanolamide, and lauric monoisopropanolamide.

A preferred concentration of the amide surfactants in the composition is 5-20% of the total active surfactant by weight.

A more preferred range is 9-16% by weight.

Amine oxide

Amine oxides useful in the present invention include long-chain alkyl amine oxides, i.e., those compounds having the formula



5 wherein R^3 is selected from an alkyl, hydroxyalkyl, acylamidopropyl and alkyl phenyl group, or mixtures thereof, containing from 8 to 26 carbon atoms, preferably 8 to 16 carbon atoms; R^4 is an alkylene or hydroxyalkylene group containing
 10 from 2 to 3 carbon atoms, preferably 2 carbon atoms, or mixtures thereof; x is from 0 to 3, preferably 0; and each R^5 is an alkyl or hydroxyalkyl group containing from 1 to 3, preferably from 1 to 2 carbon atoms, or a polyethylene oxide group containing from 1 to 3, preferably 1, ethylene oxide
 15 groups. The R^5 groups can be attached to each other, e.g., through an oxygen or nitrogen atom, to form a ring structure.

These amine oxide surfactants in particular include $\text{C}_{10}\text{-C}_{18}$ alkyl dimethyl amine oxides and $\text{C}_8\text{-C}_{12}$ alkoxy ethyl dihydroxyethyl amine oxides. Examples of such materials
 20 include dimethyloctylamine oxide, diethyldecylamine oxide, bis-(2-hydroxyethyl)dodecylamine oxide, dimethyldodecylamine oxide, dodecylamidopropyl dimethylamine oxide and dimethyl-2-hydroxyoctadecylamine oxide. Preferred are $\text{C}_{10}\text{-C}_{18}$ alkyl dimethylamine oxide, and $\text{C}_{10}\text{-C}_{18}$ acylamido alkyl dimethylamine
 25 oxide.

Most preferred amine oxides are $\text{C}_{12\text{-}14}$ alkyl dimethyl amine oxide, $\text{C}_{12\text{-}18}$ alkyl amido propyl dimethyl amine oxide, $\text{C}_{14\text{-}16}$ alkyl dimethyl amine oxide, and $\text{C}_{16\text{-}18}$ alkyl dimethyl amine oxide.

A preferred concentration for the amine oxide, or a
 30 mixture of amine oxide and betaine, in the composition is 15-60% by weight based on the total weight of active surfactant. A more preferred concentration is from 20-55% by weight. A

most preferred is 22-35% by weight.

Betaine


The betaines useful in the present invention are those compounds having the formula $R(R^1)_2N^+R^2COO^-$ wherein R is a C_6 - C_{18} hydrocarbyl group, preferably C_{10} - C_{16} alkyl group, each R^1 is typically C_1 - C_3 alkyl, preferably methyl, and R^2 is a C_1 - C_5 hydrocarbyl group, preferably a C_1 - C_5 alkylene group, more preferably a C_1 - C_2 alkylene group. Examples of suitable betaines include coconut acylamidopropyldimethyl betaine; 5 hexadecyl dimethyl betaine; C_{12} - C_{14} acylamidopropylbetaine; C_8 - C_{14} acylamidoheptyldiethyl betaine; 4- $[C_{14}$ - C_{16} acylmethylamido-diethylammonio]-1-carboxybutane; C_{16} - C_{18} acylamidodimethyl-betaine; C_{12} - C_{16} acylamidopentanedithiethyl-betaine; C_{12} - C_{16} acyl-methylamidodimethylbetaine. Preferred betaines are C_{12} - C_{18} 15 dimethylamoniohexanoate and the C_{12} - C_{18} acylamidopropane (or ethane) dimethyl (or diethyl) betaines. Also included are sulfobetaines (sultaines) of formula $R(R_1)_2N^+R_2SO_3^-$, wherein R is a C_6 - C_{18} Hydrocarbyl group, preferably a C_{10} - C_{16} alkyl group, more preferably a C_{12} - C_{14} alkyl group; each R_1 is typically C_1 - C_3 20 alkyl, preferably methyl and R_2 is a C_1 - C_6 hydrocabyl group, preferably a C_1 - C_3 alkylene or, preferably, hydroxyalkylene group. Examples of suitable sultaines are C_{12} - C_{14} di-hydroxyethylammonio propane sulfonate, and C_{16} - C_{18} dimethyl-ammonio hexane sulfonate, with C_{12} - C_{14} amido propyl ammonio-2- 25 hydroxypropyl sultaine being preferred. In a most preferred embodiment of the invention, the betaine is C_{12} - C_{18} amidopropyl betaine.

A preferred range of concentration for betaines, or mixture of betaine and amine oxide, in the composition is 15- 30 60% of the total active surfactant by weight. A more preferred range is 20-55% by weight. A most preferred range is 22-35% by weight.

Polyhydroxy fatty acid amide

The polyhydroxy fatty acid amides useful in the inventive

detergent compositions have the formula:



wherein: R¹ is H, C₁-C₄ hydrocarbyl, 2-hydroxy ethyl, 2-hydroxy propyl, or a mixture thereof, preferably C₁-C₄ alkyl, more preferably C₁ or C₂ alkyl, most preferably C₁ alkyl (i.e., methyl); and R² is a C₆-C₃₁ hydrocarbyl, preferably straight-chain C₇-C₁₉ alkyl or alkenyl, more preferably straight-chain C₉-C₁₇ alkyl or alkenyl, most preferably straight-chain C₁₁-C₁₇ alkyl or alkenyl, or mixture thereof; and Z is a polyhydroxyhydrocarbyl having a linear hydrocarbyl chain with at least 3 hydroxyls directly connected to the chain, or an alkylated derivative (preferably ethoxylated or propoxylated) thereof. Z preferably will be derived from a reducing sugar in a reductive amination reaction; more preferably Z is a glycityl. Suitable reducing sugars include glucose, fructose, maltose, lactose, galactose, mannose, and xylose. As raw materials, high dextrose corn syrup, high fructose corn syrup, and high maltose corn syrup can be utilized as well as the individual sugars listed above. These corn syrups may yield a mix of sugar components for Z. It should be understood that it is by no means intended to exclude other suitable raw materials. Z preferably will be selected from the group consisting of of -CH₂-(CHOH)_n-CH₂OH, -CH(CH₂OH)-(CHOH)_{n-1}-CH₂OH, -CH₂-(CHOH)₂(CHOR')-CH₂OH, where n is an integer from 3 to 5, inclusive, and R¹ is H or a cyclic or aliphatic monosaccharide, and alkoxyated derivatives thereof. Most preferred are glycityls wherein n is 4, particularly -CH₂-(CHOH)₄-CH₂OH.

R¹ can be, for example, N-methyl, N-ethyl, N-propyl, N-isopropyl, N-butyl, N-2-hydroxy ethyl, or N-2-hydroxy propyl.

R²-CO-N< can be, for example, cocamide, stearamide, oleamide, lauramide, myristamide, capricamide, palmitamide,

tallowamide, etc. Z can be 1-deoxyglucityl, 2-deoxyfructityl, 1-deoxymaltityl, 1-deoxylactityl, 1-deoxygalactityl, 1-deoxymannityl, 1-deoxymaltotriosityl, etc.

Alkylpolysaccharides

5 Alkylpolysaccharides such as those disclosed in U.S. Patent 4,565,647 are nonionic surfactants useful in the present invention. Suitable alkylpolysaccharides include those having a hydrophobic group containing from about 6 to about 30 carbon atoms, preferably from about 10 to about 16 carbon atoms and a
10 polysaccharide, e.g., a polyglucoside, hydrophilic group containing from about 1.3 to about 10, preferably from about 1.3 to about 3, most preferably from about 1.3 to about 2.7 saccharide units. Any reducing saccharide containing 5 or 6 carbon atoms can be used, e.g., glucose, galactose and
15 galactosyl moieties can be substituted for the glucosyl moieties. (Optionally the hydrophobic group is attached at the 2-, 3-, 4-, etc. positions thus giving a glucose or galactose as opposed to a glucoside or galactoside.) The intersaccharide bonds can be, e.g., between the one position of the additional
20 saccharide units and the 2-, 3-, 4-, and/or 6- positions on the preceding saccharide units.

As noted above, the inventive formulations include a first nonionic surfactant member selected from the group consisting of linear alcohol ethoxylates, alkyl polysaccharides, betaines,
25 and polyhydroxy fatty acid amides.


In preferred compositions, the first nonionic surfactant member is preferably a linear alcohol ethoxylate having 6-10 carbon atoms and 3 to 12 moles, more preferably 5-7 moles, of ethoxylation per mole of alcohol. Other preferred alcohol
30 ethoxylates are condensation products of aliphatic alcohols with from about 1 to about 25 moles of ethylene oxide. The alkyl chain of the aliphatic alcohol can either be straight or branched, primary or secondary, and generally contains from 8 to 22 carbon atoms. Particularly preferred are the

condensation products of alcohols having an alkyl group containing from about 10 to about 20 carbon atoms with from about 2 to about 10 moles of ethylene oxide per mole of alcohol.

5 Ethoxylated alcohols having no more than 12 carbon atoms in the alkyl position are commercially available and include NeodolJ 1-4; NeodolJ 1-7; NeodolJ 91-8, each of which is marketed by Shell Chemical Company.

10 The compositions of the invention optionally include a second nonionic surfactant member. In preferred compositions, the second member is an amine oxide in an amount of from about 1-11, more preferably 5-11, % by weight of the composition.

The preferred amine oxides for use as the second surfactant member may be represented by the general formula:
 15 $R_1R_2R_3N^+O^-$ wherein R_1 is a higher alkyl radical having from 8 to 18 carbon atoms, such as lauryl, decyl, cetyl, oleyl, stearyl, hexadecyl or an amide substituted group, such as $RCONH(CH_3)_n$, wherein RCO is a long chain alkanoyl radical and n is a small whole number; R_2 and R_3 are each lower alkyl radicals such as
 20 methyl, ethyl, propyl or a substituted lower alkyl radical such as a hydroxyethyl, hydroxyethoxyethyl, hydroxy polyethoxyethyl, etc. Examples of suitable tertiary amine oxides include lauryl dimethyl amide oxide, coconut dimethylamine oxide, dodecyl dimethyl amine oxide,



and the like.


The preferred amides are C_8 - C_{20} alkanol amides, monoethanolamides, diethanolamides, and isopropanolamides. A particularly preferred amide is a mixture of myristic
 30 monoethanolamide and lauric monoethanolamide. This preferred amide is sold by Stepan Company, Northfield, Illinois as Ninol LMP.

Cationic ammonium compound

The cationic compound is selected according to the desired end use for the formulation - typically, the cationic compound is present in amounts ranging from about 0.5 to 20% by weight of the formulation. The cationic ammonium compound is normally selected from the group consisting of quaternary ammonium salts and amine salts (salts of primary, secondary and tertiary amines).

10 1. Disinfectant formulations

In the antimicrobial or disinfectant formulations, the purpose of the quaternary ammonium disinfectants is to reduce the rate of reproduction of or kill on contact gram positive and gram negative organisms the organisms encountered in kitchen environments. Useful such disinfectants include BTC 8358 which is N-alkyl (50% C₁₄, 40% C₁₂, and 10% C₁₆) dimethyl benzyl ammonium chloride. Other quaternary ammonium salts may be any of the well-known class of quaternary ammonium germicides characterized by the formula:



20 wherein at least one of the radicals, R₁, R₂, R₃, and R₄ ("the 'R' groups") is a hydrophobic, aliphatic, aryl aliphatic, or aliphatic aryl radical of from 6 to 26 carbon atoms, the entire cation portion of the molecule has a molecular weight of at least 165, and the remaining R groups are hydrophobic, aliphatic, aryl aliphatic, or aliphatic aryl radical of from 6 to 26 carbon atoms. The hydrophobic radicals may be long-chain alkyl, long-chain alkoxy aryl, long-chain alkyl aryl, halogen-substituted long-chain alkyl aryl, long-chain alkyl phenoxy alkyl, aryl alkyl, and so forth, in nature. The remaining radicals on the nitrogen atom other than the hydrophobic

radicals are substituents of hydrocarbon structure usually containing a total of no more than 12 carbon atoms. The radical X in the above formula is any salt-forming anionic radical.


- 5 Suitable quaternary ammonium salts within the above description include the alkyl ammonium halides such as cetyl trimethyl ammonium bromide, alkyl aryl ammonium halides such as octadecyl dimethyl benzyl ammonium bromide, N-alkyl pyridinium halides such as N-cetyl pyridinium bromide, and the like.
- 10 Other suitable types of quaternary ammonium salts include those in which the molecule contains either, amide or ester linkages such as octyl phenoxy ethoxy ethyl dimethyl benzyl ammonium chloride. N-(laurylcocoaminoformylmethyl) - pyridinium chloride, and so forth. Other very effective types of
- 15 quaternary ammonium germicides are those in which the hydrophobic radical is characterized by a substituted aromatic nucleus as in the case of lauryloxyphenyltrimethyl ammonium chloride, cetylaminophenyltrimethyl ammonium methosulfate, dodecylphenyl-trimethyl ammonium methosulfate,
- 20 dodecylbenzyltrimethyl ammonium chloride, chlorinated dodecylbenzyltrimethyl ammonium chloride, and the like.

Preferred quaternary ammonium germicides of the above general types are the long-chain alkyl dimethylbenzyl quaternary ammonium salts, the alkyl phenoxy alkoxy alkyl

25 dimethyl benzyl quaternary ammonium salts, the N-(acylcocoaminoformylmethyl)-pyridinium halides, the long-chain alkyl trimethyl ammonium halides, the long-chain alkyl benzyl dimethyl benzyl ammonium halides, and the long-chain alkyl benzyl diethyl ethanol ammonium halides in which the alkyl

30 radical contains from 8-18 carbon atoms.

The quaternary ammonium salts useful in the invention have the general formula:



wherein R_1 and R_2 are straight or branched chain lower alkyl groups having from one to seven carbon atoms; R_3 is a straight or branched chain higher alkyl group having from about six to
 5 sixteen carbon atoms, or a benzyl group; R_4 is a straight or branched chain higher alkyl group having from about six to sixteen carbon atoms; and X is a halogen or a methosulfate or saccharinate ion.

In preferred quaternary ammonium salts, R_1 and R_2 are
 10 methyl groups; R_3 is benzyl or straight or branched chain alkyl having from about eight to sixteen carbon atoms; and R_4 is straight or branched chain alkyl having from about eight to sixteen carbon atoms provided that not both R_3 and R_4 have
 15 sixteen carbon atoms simultaneously. A preferred halogen is chlorine, or a methosulfate or a saccharinate ion.

Illustrative of suitable quaternary ammonium germicides are: dioctyl dimethyl ammonium chloride, octyl decyl dimethyl ammonium chloride, didecyl dimethyl ammonium chloride, (C_{12} - C_{18}) n-alkyl dimethyl benzyl ammonium chloride, (C_{12} - C_{18}) n-alkyl
 20 dimethyl ethylbenzyl ammonium chloride, (C_{12} - C_{18}) n-alkyl dimethyl benzyl ammonium saccharinate, di(C_1 - C_7) alkyl (C_6 - C_{26}) alkyl aryl ammonium salts, di(C_1 - C_7)alkyl di(C_6 - C_{14}) alkyl ammonium salts, tri(C_1 - C_7) alkyl (C_6 - C_{26}) alkyl ammonium salts, (C_{14} - C_{26})alkyl di(C_1 - C_7)alkyl aryl ammonium salts, and (C_{14} -
 25 C_{16})alkyl tri (C_1 - C_7) alkyl aryl ammonium salts. This is not an exhaustive list and other quaternary ammonium salts having germicidal activity will suffice. The quaternary ammonium salt in the present invention need not be a single entity, but may be a blend of two or more quaternary ammonium salts.

30 The more preferred quaternary salts are alkyl trimethyl ammonium compounds and alkyl dimethyl benzyl (and/or

ethylbenzyl) ammonium compounds having from 10-18 carbon atoms in the long chain portion. The most preferred quaternary salts are Benzalkonium (C_{12-16}) chloride, alkyl (cetyl or oleyl) trimethyl ammonium chloride, or other biocidal quats such as
5 dialkyl dimethyl quats.

Where the cationic surfactant is a dialkyl dimethyl ammonium compound, the carbon chains should have from 6-14 carbons atoms in each of the two long chain alkyl groups. Di(C_{16-18}) alkyl ammonium compounds are most preferably avoided
10 in the compositions and formulations of the invention.

The amount, in weight-percent, of the quaternary ammonium salt, either as a single entity or blended, is up to about 50%, but is preferably from about 0.1%-10.0% and more preferably about 1-3%. The preferred quaternary ammonium germicide is a
15 mixture of about 34% by weight C_{12} and 16% by weight C_{14} n-alkyl dimethyl ethylbenzyl ammonium chloride and about 30% by weight C_{14} , 15% by weight C_{16} , 2.5% by weight C_{12} and 2.5% by weight C_{18} n-alkyl dimethyl benzyl ammonium chloride.

The quaternary ammonium compounds can also function as
20 cationic surfactants to produce antistatic and conditioning effects when deposited on the substrate.

2. Conditioning formulations

The invention also encompasses cleaning compositions capable of imparting a conditioning effect on a substrate such
25 as skin or hair. Such formulations include hand soaps and conditioning shampoos. A variety of cationic surfactants useful as deterative surfactants and as conditioning agents are well known in the art. These materials contain amino or quaternary ammonium hydrophilic moieties which are positively
30 charged when dissolved in the aqueous composition of the present invention. Whether the cationic surfactant functions as a deterative surfactant or a conditioning agent, or both, will depend upon the particular compound as is well understood by those skilled in the art. In general, compounds with longer

chain length moieties attached to the cationic nitrogen tend to exhibit greater conditioning benefits. Cationic surfactants among those useful herein are disclosed in the following documents, all incorporated by reference herein: M.C. Publishing Co., *McCutcheon's, Detergents & Emulsifiers*, (North American edition 1993); Schwartz et al., *Surface Active Agents, Their Chemistry and Technology*, New York; Interscience Publishers, 1949; U.S. Patent 3,155,591, Hilfer, issued November 3, 1964; U.S. Patent 3,929,678, Laughlin et al.,
10 issued December 30, 1975; U.S. Patent 3,959,461, Bailey et al., issued May 25, 1976; and U.S. Patent 4,387,090, Bolich, Jr., issued June 7, 1983.

Quaternary ammonium salts include dialkyl dimethyl-ammonium chlorides and trialkyl methyl ammonium chlorides, wherein the
15 alkyl groups have from about 12 to about 22 carbon atoms and are derived from long-chain fatty acids. These types of cationic surfactants are useful as hair conditioning agents. Examples of quaternary ammonium salts useful herein include di(coconutalkyl) dimethyl ammonium chloride, stearyl dimethyl
20 benzyl ammonium chloride. Stearyl dimethyl benzyl ammonium chloride and cetyl trimethyl ammonium chloride are preferred quaternary ammonium salts useful herein. (Hydrogenated tallow) trimethyl ammonium chloride is a preferred quaternary ammonium salt. Preferred of the conventional cationic conditioning
25 agents are cetyl trimethyl ammonium chloride, lauryl trimethyl ammonium chloride, stearyl dimethyl benzyl ammonium chloride, and (partially hydrogenated tallow) trimethyl ammonium chloride; these materials may also provide anti-static benefits to the present shampoo compositions.

30 Salts of primary, secondary and tertiary fatty amines are also suitable cationic surfactant materials. The alkyl groups of such amines preferably have from about 12 to about 22 carbon atoms, and may be substituted or unsubstituted. Secondary and tertiary amines are preferred, tertiary amines are particularly

preferred. Such amines, useful herein, include stearamido propyl dimethyl amine, diethyl amino ethyl stearamide, dimethyl stearamine, dimethyl soyamine, soyamine, myristyl amine, tridecyl amine ethyl stearylamine, N-tallowpropane diamine, 5 ethoxylated (5 moles E.O.) stearylamine, dihydroxy ethyl stearylamine, and arachidylbehenylamine. Suitable amine salts include the halogen, acetate, phosphate, nitrate, citrate, lactate and alkyl sulfate salts. Such salts include stearylamine hydrochloride, soyamine chloride, stearylamine 10 formate, N-tallowpropane diamine dichloride and stearamidopropyl dimethylamine citrate. Cationic amine surfactants included among those useful in the present invention are disclosed in U.S. Patent 4,275,055, Nachtigal, et al., issued June 23, 1981, incorporated by reference herein.

15 Cationic conditioning surfactants especially useful in shampoo formulations are quaternary ammonium or amino compounds having at least one N-radical containing one or more nonionic hydrophilic moieties selected from alkoxy, polyoxyalkylene, alkylamido, hydroxyalkyl, and alkylester moieties, and 20 combinations thereof. The surfactant contains at least one hydrophilic moiety within 4, preferably within 3, carbon atoms (inclusive) of the quaternary nitrogen or cationic amino nitrogen. Additionally, carbon atoms that are part of a hydrophilic moiety, e.g., carbon atoms in a hydrophilic 25 polyoxyalkylene (e.g., $-\text{CH}_2-\text{CH}_2-\text{O}-$), that are adjacent to other hydrophilic moieties are not counted when determining the number of hydrophilic moieties within 4, or preferably 3, carbon atoms of the cationic nitrogen. In general, the alkyl portion of any hydrophilic moiety is preferably a C_1 - C_3 alkyl. 30 Suitable hydrophile-containing radicals include, for example, ethoxy, propoxy, polyoxyethylene, polyoxypropylene, ethylamido, propylamido, hydroxymethyl, hydroxyethyl, hydroxypropyl, methylester, ethylester, propylester, or mixtures thereof, as nonionic hydrophile moieties. The amino surfactants must be

positively charged at the pH of the shampoo compositions. Generally, the pH of the shampoo compositions will be less than about 10, typically from about 3 to about 9.

Other cationic compounds suitable for use in the invention include NH_4^+ , and mono-, di-, and tri-short chain alkyl ammonium salts.

Adjuvant materials

Various adjuvant materials may be added to these foaming aqueous detergent compositions such as small amounts of viscosity builders and conditioning agents inclusive of gums and hydroxy-propyl methyl cellulose. Such ingredients can be added in an amount that does not adversely effect the foaming and cleaning characteristics of the compositions. Other ingredients may include alkaline or acid buffers to aid in the adjustment and maintenance of the desired pH of the finished product such as borax, various inorganic water-soluble phosphates, sodium hydroxide, citric acid, etc. Other additions include optical brighteners, bleaches, germicides, fungicides, bactericides, colorants, perfumes, etc. in minor amounts which do not interfere with the cleaning, foaming, conditioning, or sanitizing properties of the composition.

Other ingredients include ethylenediamine tetraacetate (EDTA) and polyethylene glycol (PEG) fatty acid esters. EDTA is especially useful in antibacterial cleaning compositions, particularly handsoaps, since it increases the effectiveness of the antibacterial quaternary ammonium compound, in particular against *Pseudomonas Aeruginosa*, a pathogenic gram-negative organism. A representative PEG ester is PEG 600 distearate. This ester provides excellent viscosity enhancement in the inventive surfactant systems by association with other components without causing a loss of clarity or an increase in color.

All documents, e.g., patents and journal articles, cited above or below are hereby incorporated by reference in their

entirety.

In the examples, all amounts are stated in percent by weight of active material unless indicated otherwise.

One skilled in the art will recognize that modifications
5 may be made in the present invention without deviating from the spirit or scope of the invention. The invention is illustrated further by the following examples which are not to be construed as limiting the invention or scope of the specific procedures or compositions described herein.

10 The detergent compositions of the invention are prepared by combining water with the alcohol ethoxylate and amine oxide surfactant members, mixing until uniform and then adding any optional components, such as, for example, amide or cationic ammonium compound, and again mixing until uniform. Heating may
15 be employed as needed to enhance dissolution of the components.

Various antimicrobial cleaning formulations described herein were analyzed for detergency and foam height. Detergency and foam longevity was evaluated using the miniplate assay described below. Foam height was evaluated according to
20 the Ross Miles Foam Height test as described by J. Ross and J.D. Miles in Oil and Soap, 18; 99 (1941) at 0.032% active concentration using water of 140 ppm hardness at 25°C. Ross-Miles test results are displayed in cm.

Foam longevity and Detergency Evaluation

25 The capability of various formulations for cleaning and degreasing was determined by the Mini-Plate Test, as follows:

Preparation of Soil Material

1. Melt shortening (Crisco, approx. 100 g) in a beaker at 160°F.
- 30 2. Add a small amount (not much needed for deep color) of red dye to melted Crisco and stir until dissolved.
3. Calibrate syringe to deliver 0.36 g of Crisco soil on each plate.
4. Apply 0.36 g of Crisco oil to each of the watchglasses

- (One large watchglass is equivalent to three mini-plates).
5. When all of the larger watchglasses have been soiled, recalibrate syringe to deliver 0.12 g of Crisco soil to each plate.
 - 5 6. Apply 0.12 g of Crisco soil to each of the smaller watchglasses.
 7. Allow soiled watchglasses to harden at room temperature overnight before using.
 8. Soiled watchglasses should always be stored at room
10 temperature [(can be stored indefinitely)].

Procedure for Analyzing Test Formulations

1. A test solution may be made by diluting sufficient product with [tap] water (140 ppm hardness) to a concentration of
15 0.048% active material. 400 ml of such a solution is employed; heating to about 130-135°F may be necessary to achieve dissolution of the product.
3. The solution is placed in a Pyrex dish and then agitated with a paintbrush to generate foam, and the temperature of
20 the solution adjusted to 120°F.
4. At this point, large watchglasses (which represent three plates each) are washed, one every 45 seconds, by removing a thin layer of soil at a time from the surface of the plate with the paintbrush, then agitating the paintbrush
25 in the solution to remove the adhering soil (which consequently breaks down the foam).
5. The endpoint of the test is the number of mini-plates washed when further agitation of the solution fails to produce new/additional foam on the solution surface. In
30 certain tests, small watchglasses representing 1/3 the surface area of a large watchglass may be used as the endpoint is neared.

Formulations according to the invention are shown below in

the following examples. In all the examples, all amounts are given in percent active by weight of the final formulation unless indicated to the contrary.

	FORMULATION NO.					
	1	2	3	4	5	6
C ₆₋₁₀ alkyl ethoxylate (4EO ¹) (commercially available from Stepan Company as Bio- soft FF-600)	16.50	16.50	14.00	17.60	14.85	13.20
Cocamido propyl dimethyl amine oxide (C ₁₂₋₁₈ alkyl) (commercially available from Stepan Company as Ammonyx CDO)	8.50	8.50	10.50	9.06	7.65	6.80
Lauric (C ₁₂) diethanol amide (commercially available from Stepan Company as Ninol 96-SL)	2.50	1.00	2.50	2.70	2.25	2.00
Lauric (C ₁₂) mono ethanol amide (commercially available from Stepan Company as Ninol LMP (C12))	2.50	4.00	2.50	2.70	2.25	2.00
Benzalkonium (C ₁₂₋₁₆) chloride (commercially available from Stepan Company as BTC-835)	2.00	2.00	2.00	0.00	5.00	8.00
Mini-plates	38	44	44	42	38	35
Ross-Miles, cm	14.00	11.75	14.00	12.50	14.85	14.90

¹ EO refers to the average number of moles of ethylene oxide per mole of alcohol.

	7	8	9	10	11	12	13	14
C ₆₋₁₀ alkyl ethoxylate (4EO) (Bio-soft FF-600)		8.50	11.00	9.36		9	9	16.5
Cocamido propyl dimethyl amine oxide (C ₁₂₋₁₈ alkyl)	10	16.50	5.66	4.82	4.5	23		
Alkyl polyglycoside (Glucopon 600-CS)	5							8.5
Lauric (C ₁₂) diethanol amide	2.5	2.50	1.67	1.41	2.0			2.5
Lauric (C ₁₂) Myristic (C ₁₄) mono ethanol amide	2.5	2.50	1.67	1.41	2.0			2.5
Benzalkonium (C ₁₂₋₁₆) chloride	2.0	2.00	12.00	15.00	2.0			2.0
Lonzaine JS (sulfobetaine)	16.5							
Cocamidopropyl Betaine (Amphosol CA)					4.5		23	
C ₉₋₁₁ alkyl ethoxylate (8EO)					8.5			
cetyltrimethyl ammonium chloride (commercially available from Stepan Company as Ammonyx Cetac)					1.5			
tallow amine-2EO					2.0			
NH ₄ Cl (aq.)					4.0			
Mini-plates	39	46	32	21	26	34	33	33
Ross-Miles, cm	12.75	13.81	14.10	13.90	13.4	15.15	12.6	11

	15	16	17
C ₆₋₁₀ alkyl ethoxylate (4EO)			14.8
C ₁₁ alkyl ethoxylate (7EO)	16.50		
C ₉₋₁₁ alkyl ethoxylate (8EO)		16.50	
Cocamido propyl dimethyl amine oxide (C ₁₂₋₁₃ alkyl)	8.50	8.50	7.65
Lauric (C ₁₂₋₁₄) diethanol amide	2.50	2.50	2.20
Lauric (C ₁₂) Myristic (C ₁₄) monoethanol amide	2.50	2.50	2.20
Benzalkonium (C ₁₂₋₁₆) chloride	2.00	2.00	2.00

CHEMICAL NAME	TRADE NAME	% actives, w/w	
		18	19
Alcohol Ethoxylate (C ₈₋₁₀ 9EO)	Alfonic 810-9	5.30	6
Alcohol Ethoxylate (C ₈₋₁₀ 12EO)	Alfonic 810-12	16.79	19
Cocoamidopropylamine Oxide	Ammonyx CDO	10.59	12
Lauric Diethanolamide	Ninol 96-SL	2.65	3.0
Lauric Monoethanolamide	Ninol C12LMP	2.65	3.0
Alkyl Dimethyl Benzyl Ammonium Chloride	BTC 8358	2.0	2.0
Xanthan Gum	Kelzan T, K5C487	0.07	-
Deionized Water		Q.S. to 100	Q.S. to 100

Formulation 47 can be made at 47% actives utilizing the same component ratios.

Broad Spectrum Antimicrobial Hand Soap Formulation

	Formulation No. (% actives, by weight)					
	20	21	22	23	24	25
Deionized Water	Q.S.to 100	Q.S.to 100	Q.S.to 100	Q.S.to 100	Q.S.to 100	Q.S.to 100
C ₆₋₁₀ alkyl ethoxylate (4EO) (Bio-soft FF- 600)	1.70	1.70	1.70	1.70	0-10%	0-10%
Amphosol CA (Cocamidopropyl Betaine)	5.00	16.67	5.00	16.67	0-20%	0-66%
Ammonyx CDO (Cocamidopropyl amine Oxide)	2.50	8.33	2.50	8.33	0-10%	0-33%
Ammonyx CO (Cocamine Oxide)	1.00	3.33	1.00	3.33	0-10%	0-33%
Kessco PEG 6000 DS (PEG 6000 Distearate)	-	-	1.00	1.00	0-5%	0-5%
Ninol LMP (Lauramide MEA)	2.00	2.00	2.00	2.00	0-10%	0-10%
BTC 835 (Benzalkonium Chloride)	1.00	2.00	1.00	2.00	0-5%	0-10%
Dow Versene 100 (Tetrasodium EDTA)	0.20	0.51	0.20	0.51	0-0.5%	0-1.3%
NH ₄ Cl (Optional)	4.00	4.00	-	-	-	-
Viscosity (cps)		1920		7290		
Ross Miles, cm		12.8		8.2		
Mini-Plates		39		42		

5

	FORMULATION NO.					
	26	27	28	29	30	31
C ₆₋₁₀ alkyl ethoxylate (4EO) (Bio-soft FF-600)	16.50					
Cocamido propyl dimethyl amine oxide (C ₁₂₋₁₈ alkyl) (Ammonyx CDO)	8.50	8.50	8.50	8.50	8.50	8.50
Lauric (C ₁₂) diethanol amide (Ninol 96-SL)	2.50	2.50	2.50	2.50	2.50	2.50
Lauric (C ₁₂) Myristic (C ₁₄) mono ethanol amide (Ninol LMP (C12))	2.50	2.50	2.50	2.50	2.50	2.50
Benzalkonium (C ₁₂₋₁₆) chloride (BTC-835)	2.00	2.00	2.00	2.00	2.00	2.00
Linear C ₁₀₋₁₂ alkyl (6EO) ethoxylate (BioSoft ET-630)		16.50				
Linear primary alcohol ethoxylates (3EO) (Neodol 1-3)			16.50			
Linear primary alcohol ethoxylates (5EO) (Neodol 1-5)				16.50		
Linear primary alcohol ethoxylates (7EO) (Neodol 1-73B)					16.50	
Alkylpolyglycoside (C ₁₂₋₁₄ alkyl) (Glucopon 625)						16.50
Mini-Plates	38	36	38	33	40	25
Ross Miles, cm	12.70	11.50	2.50	9.50	9.50	9.50

5

	FORMULATION NO.					
	32	33	34	35	36	37
C ₆₋₁₀ alkyl ethoxylate (4EO) (Bio-soft FF-600)		16.50			16.50	
Cocamido propyl dimethyl amine oxide (C ₁₂₋₁₈ alkyl) (Ammonyx CDO)	8.50			8.50		
Lauric (C ₁₂) diethanol amide (Ninol 96-SL)	2.50	2.50	2.50	2.50	2.50	2.50
Lauric (C ₁₂) Myristic (C ₁₄) mono ethanol amide (Ninol LMP (C12))	2.50	2.50	2.50	2.50	2.50	2.50
Benzalkonium (C ₁₂₋₁₆) chloride (BTC-835)	2.00	2.00	2.00	2.00	2.00	2.00
Glucosamide (C ₁₂₋₁₄)	16.50	8.50				
Linear primary alcohol ethoxylates (C ₉₋₁₁ alkyl) (Neodol 91-8)			16.50	16.50		16.50
Laurylamine oxide (Ammonyx LO)			8.50			
Myristylamine oxide (Ammonyx MO)					8.50	8.50
Mini-Plates	36		38	37	41	37
Ross Miles, cm	2.00	9.25	9.50	10.20	12.00	11.20

5

	FORMULATION NO.				
	38	39	40	41	42
C ₆₋₁₀ alkyl ethoxylate (4EO) (Bio-soft FF-600)	16.50	16.50	16.50	16.50	16.50
Cocamido propyl dimethyl amine oxide (C ₁₂₋₁₈ alkyl) (Ammonyx CDO)	8.50	8.50	8.50		
Lauric (C ₁₂) diethanol amide (Ninol 96-SL)	2.50	2.50	2.50	2.50	2.50
Lauric (C ₁₂) Myristic (C ₁₄) mono ethanol amide (Ninol LMP (C12))	2.50	2.50	2.50	2.50	2.50
Benzalkonium (C ₁₂₋₁₆) chloride (BTC-835)				2.00	2.00
Cetyltrimethyl ammonium chloride (Ammonyx Cetac)		2.00			
Olealkonium chloride (Ammonyx KP)			2.00		
Cocamidopropyl Betaine (Amphosol CG)					8.50
Alkylpolyglycoside (C ₁₂₋₁₄ alkyl) (Glucopon 625)				8.50	
pH	8.58	8.50	8.50	9.70	8.60
Mini-Plates	32	32	35	33	33
Ross Miles, cm	10.55	16.00	13.80	11.00	15.00

5

	FORMULATION NO.					
	43*	44	45	46	47	48
C ₆₋₁₀ alkyl ethoxylate (4EO) (Biosoft FF-600)		16.50	22.50	17.33	17.03	16.83
Lauric (C ₁₂₋₁₄) diethanol amide (Ninol 96-SL)	2.50	2.50	3.40	2.63	2.58	2.55
Lauric (C ₁₂) Myristic (C ₁₄) mono ethanol amide (Ninol LMP (C12))	2.50	2.50	3.40	2.63	2.58	2.55
Benzalkonium (C ₁₂₋₁₆) chloride (BTC-835)	2.00	2.00	2.72			
Quarternium 24 (and) Benzalkonium chloride (BTC 885)				0.53	1.03	1.53
Cocamidopropyl Betaine (Amphosol CG)	8.50					
Alkylpolyglycoside (C ₁₂₋₁₄ alkyl) (Glucopon 625)	16.50					
Mini-Plates	23	29	30	38	38	37
Ross Miles, cm	11.10	10.10	11.00	9.60	10.60	11.90

10

*Comparative example

5

	FORMULATION NO.					
	49	50	51	52	53	54
C ₆₋₁₀ alkyl ethoxylate (4EO) (Bio-soft FF-600)	16.50	16.50	16.50	16.50	16.50	16.50
Cocamido propyl dimethyl amine oxide (C ₁₂₋₁₈ alkyl) (Ammonyx CDO)	8.50	8.50	8.50	8.50	8.50	8.50
Lauric (C ₁₂) diethanol amide (Ninol 96-SL)	5.00					
Lauric (C ₁₂) mono ethanol amide (Ninol LMP (C12))		5.00				
Benzalkonium (C ₁₂₋₁₆) chloride (BTC-835)	2.00	2.00	2.00	2.00	2.00	2.00
Coco diethanolamide (Ninol 40-CO)			5.00			
Lauric/Myristic monoethanolamide (Ninol LMP)				5.00		
Coco monoisoproanolamide (Ninol M-10)					5.00	
Coco diethanolamide, modified (Ninol 11-CM)						5.00
Mini-Plates	33	41	33	39	36	36
Ross Miles, cm	13.70	14.00	10.50	13.10	12.00	9.00

5

	FORMULATION NO.					
	55	56	57	58	59	60
C ₆₋₁₀ alkyl ethoxylate (4EO) (Bio-soft FF-600)	16.50	16.50	16.50			
Cocamido propyl dimethyl amine oxide (C ₁₂₋₁₈ alkyl) (Ammonyx CDO)	8.50			4.25	8.50	8.50
Lauric (C ₁₂) diethanol amide (Ninol 96-SL)	2.50	5.00	2.50	2.50	2.50	2.50
Myristylamine oxide (Ammonyx MO)		8.50				
Benzalkonium (C ₁₂₋₁₆) chloride (BTC 835)	2.00			2.00	2.00	2.00
Myristic/Cetyl dimethylamine oxide (Ammonyx MCO special)				4.25		
Lauric/Myristic monoethanolamide (Ninol LMP)	2.50		2.50	2.50	2.50	2.50
Stearamine oxide (Ammonyx SO)			8.50			
Linear primary alcohol ethoxylates (C ₉₋₁₁ alkyl) (Neodol 91-8)				16.50	16.50	16.50
Quarternium 12 (BTC 1010)		2.00	2.00			
Xanthan gum (Keltrol RD)					0.01	
Methocel J40MS-N						0.10
Mini-Plates	37	41	37			
Ross Miles, cm	14.50	9.50	14.30	11.5		9.05

5

	FORMULATION NO.					
	61	62	63	64	65	66
C ₆₋₁₀ alkyl ethoxylate (4EO) (Bio-soft FF-600)		16.50	16.50	16.50		
Cocamido propyl dimethyl amine oxide (C ₁₂₋₁₈ alkyl) (Ammonyx CDO)	8.50	8.50				
Lauric (C ₁₂) diethanol amide Ninol 96-SL	2.50	5.00	2.50	2.50	2.50	2.50
Myristic/Cetyl dimethylamine oxide (Ammonyx MCO special)						8.50
Benzalkonium (C ₁₂₋₁₆) chloride (BTC-835)	2.00				2.00	2.00
Cetamine oxide (Ammonyx CO)				8.50	8.50	
Lauric/Myristic monoethanolamide (Ninol LMP)	2.50	2.50	2.50	2.50	2.50	2.50
Stearamine oxide (Ammonyx SO)			8.50			
Linear primary alcohol ethoxylates (C ₉₋₁₁ alkyl) (Neodol 91-8)	16.50				16.50	16.50
Quarternium 12 (BTC 1010)		2.00	2.00	2.00		
Xanthan gum (Keltrol RD)		0.01				
PEG-150 distearate (Kesco PEG 6000)	2.00					
Mini-Plates			35			
Ross Miles, cm	10.15	14.10	14.50	12.50	11.50	

5

	FORMULATION NO.					
	67	68	69	70	71	72
C ₆₋₁₀ alkyl ethoxylate (4EO) (Bio-soft FF-600)			16.50	16.50	16.50	16.50
Cocamido propyl dimethyl amine oxide (C ₁₂₋₁₈ alkyl) (Ammonyx CDO)			8.50	8.50	8.50	8.50
Lauric (C ₁₂) diethanol amide (Ninol 96-SL)	2.50	2.50	2.50	2.50	2.50	2.50
Myristamine oxide (Ammonyx MO)		8.50				
Benzalkonium (C ₁₂₋₁₆) chloride (BTC-835)	2.00	2.00				2.00
Cetamine oxide (Ammonyx CO)						
Lauric/Myristic monoethanolamide (Ninol LMP)	2.50	2.50	2.50	2.50	2.50	2.50
Stearamine oxide (Ammonyx SO)	8.50					
Linear primary alcohol ethoxylates (C ₉₋₁₁ alkyl) (Neodol 91-8)	16.50	16.50				
Quarternium 12 (BTC 1010)			2.00			
Quarternium 24 (BTC 818)				2.00		
Quarternium 24 (and) Benzalkonium chloride (BTC 885)					2.00	
Mini-Plates			40	40	37	40
Ross Miles, cm			13.95	13.45	14.08	13.65

5

FORMULATION NO.

	73	74	75	76
C ₆₋₁₀ alkyl ethoxylate (4EO) (Bio-soft FF- 600)	13.20	11.50	14.80	14.80
Cocamido propyl dimethyl amine oxide (C ₁₂₋₁₈ alkyl) (Ammonyx CDO)	6.80	5.58	7.65	7.65
Lauric (C ₁₂) diethanol amide (Ninol 96-SL)	2.00	1.80	2.20	2.20
Benzalkonium (C ₁₂₋₁₆) chloride (BTC-835)	2.00	2.00	2.00	2.00
Lauric/Myristic monoethanolamide (Ninol LMP)	2.00	1.80	2.20	2.20
Acusol ACNSO 880			1.00	2.00
Mini-Plates	33	32		
Ross Miles, cm	13.69	12.54		

5

	FORMULATION NO.					
	77	78	79	80*	81*	82
C ₆₋₁₀ alkyl ethoxylate (4EO) (Bio-soft FF-600)	16.50	16.50	16.50			
Lauric (C ₁₂) diethanol amide (Ninol 96-SL)	2.50	2.50	2.50	2.50	2.50	2.50
Myristylamine oxide (Ammonyx MO)				8.50		
Benzalkonium (C ₁₂₋₁₆) chloride (BTC-835)	2.00	2.00	2.00	2.00	2.00	2.00
Cocoamidopropylamine oxide (Ammonyx CDO)					8.50	8.50
Lauric/Myristic monoethanolamide (Ninol LMP)	2.50	2.50	2.50			
Lauric (C ₁₂) Myristic (C ₁₄) mono ethanol amide (Ninol LMP (C12))				2.50	2.50	2.50
Glucosamide (C ₁₂₋₁₄) (Mega 12/14)						16.50
sulfobetaine (Lonzaine JS)	14.15					
[chem. Name] (APG 600 CS)		4.25	8.50	16.50	16.50	
Mini-Plates			35	39	25	36
Ross Miles, cm	10.39	10.25	11.01	8.10	9.50	2.00

*Comparative example

Hand Soap and Conditioning Shampoo Formulations

Hand soap and conditioning shampoo formulations according to the invention are prepared essentially by the method described above for dishwashing compositions.

Ingredient	Formulation				
	A	B	C	D	E
C ₉₋₁₁ alkyl ethoxylate (8EO)			4.40		2.30
C ₆₋₁₀ alkyl ethoxylate (4EO)	5.50	4.40		2.30	
cocoamidopropyl dimethyl amine oxide	2.83	2.30	2.30	4.40	4.40
lauric diethanolamide	0.83			0.65	0.65
lauric myristic monoethanolamide	0.83	1.30	1.30	0.65	0.65
cetyl trimethyl ammonium chloride					1.00
benzalkonium chloride	2.00	1.00	1.00	1.00	1.00
water	q.s.	q.s.	q.s.	q.s.	q.s.
Total Actives	11.99	9.00	9.00	9.00	10.00

Comparative Examples				
	I	II	III	IV
C ₁₂₋₁₃ alkyl ethoxylate (6.5EO)	5	14		
C ₁₂₋₁₅ alkyl ethoxylate (9EO)	5			
C ₁₂ amine oxide	5			
cocoamidopropyl betaine	5	5		5
Lonzaine JS (sulfobetaine)	5			12
Lauric myristic cocodiethanolamide	5	3	0.8	
Oleyl dimethyl ammonium chloride				1.5
Cocodiethanolamide				3
Cocamido propyl dimethyl amine oxide (C ₁₂₋₁₄ alkyl)				5
Benzalkonium (C ₁₂₋₁₆) chloride	1.4	1.6	9.2	1
Mini-plates	21	24	42	31
Ross-Miles, cm	7.3	7.15	10.10	10.85

From the foregoing, it will be appreciated that although specific embodiments of the invention have been described herein
5 for purposes of illustration, various modifications may be made without deviating from the spirit or scope of the invention.

What is claimed is:

1. An aqueous liquid cleaning composition, the composition being free of anionic surfactants and comprising a nonionic surfactant system consisting essentially of:

5 (a) from about 0.1-50% by weight based on the weight of the composition of a linear alcohol ethoxylate having an average carbon chain length of no more than 12 carbon atoms; and

10 (b) a surfactant member selected from the group consisting of amine oxides, betaines, and mixtures thereof, the total concentration of active components in the composition based on the weight of the composition is at least about 5%.

2. A composition according to claim 1, further including
15 a surfactant selected from the group consisting of alkanolamides, alkyl polysaccharides, betaines, cationic ammonium compounds, polyhydroxy fatty acid amides, and mixtures thereof.

20 3. A composition according to Claim 2, comprising from about 0.5-20% by weight of the composition of a cationic ammonium compound.

4. A composition according to claim 2, wherein the amount
25 of linear alcohol ethoxylate is from about 8-18% by weight of the composition.

5. A composition according to claim 4, wherein the amount
30 of linear alcohol ethoxylate is from about 9-17% by weight of the composition.

6. A composition according to claim 4, wherein the amount of the amine oxide in the composition is from about 3-23% by

weight of the composition.

7. A composition according to claim 1, wherein the amount of linear alcohol ethoxylate has from 6-11 carbon atoms in the alkyl portion and from about 4-8 moles of ethoxylation per mole of alcohol.

8. A composition according to claim 1, wherein the amounts of linear alcohol ethoxylate and amine oxide in the composition are respectively, by weight of the composition, from about 9-18%, and 3-23%.

9. An aqueous liquid cleaning composition, the composition being free of anionic surfactants and consisting essentially of:

- a) from about 2-23% by weight based on the weight of the composition of a linear alcohol ethoxylate having an average carbon chain length of no more than 12 carbon atoms;
 - (b) from about 2-23% by weight based on the weight of the composition of a surfactant member selected from the group consisting of amine oxides, betaines, and mixtures thereof; and
 - (c) a cationic ammonium compound,
- the total concentration of active components in the composition based on the weight of the composition is at least about 5%.

10. A composition according to Claim 9, wherein the cationic ammonium compound is a quaternary ammonium disinfecting compound.

11. A composition according to Claim 10, wherein the linear alcohol ethoxylate has a minimum of 6 carbon atoms and 3

to 12 moles of ethoxylation per mole of alcohol.

12. A composition according to claim 11, wherein the amount of linear alcohol ethoxylate is from about 8-18% by weight of the composition.

13. A composition according to claim 12, wherein the amount of linear alcohol ethoxylate is from about 9-17% by weight of the composition.

14. A composition according to claim 13, wherein the amount of the amine oxide in the composition is from about 3-23% by weight of the composition.

15. A composition according to claim 14, wherein the amounts of linear alcohol ethoxylate and amine oxide in the composition are respectively, by weight of the composition, from about 9-18%, and 3-23%.

16. A method for preparing a cleaning composition comprising:

(a) preparing a surfactant system consisting essentially of:

(i) from about 0.1-50% by weight based on the weight of the composition of a linear alcohol ethoxylate having an average carbon chain length of no more than 12 carbon atoms; and

(ii) a surfactant member selected from the group consisting of amine oxides, betaines, and mixtures thereof;

and

(b) combining the surfactant system with a cationic ammonium compound such that the total concentration of active components in the composition based on the weight of the composition is at least about 5%.

17. An aqueous liquid cleaning composition, the composition being free of anionic surfactants and comprising a nonionic surfactant system consisting essentially of:

- 5 (a) from about 13-19% by weight based on the weight of the composition of a linear alcohol ethoxylate having an average carbon chain length of no more than 12 carbon atoms; and;
- (b) from about 3-7% by weight of the composition of a
10 mono- or dialkanolamide; and
- (c) from about 5-10% by weight of the composition of an alkylpolyglycoside.

18. An aqueous liquid cleaning composition, the
15 composition being free of anionic surfactants and comprising a nonionic surfactant system consisting essentially of:

- (a) from about 13-19% by weight based on the weight of the composition of a sulfobetaine;
- (b) from about 5-20% by weight of the composition of
20 an amine oxide, a betaine, or mixture thereof;
- (c) from about 3-7% by weight of the composition of a mono- or dialkanolamide or mixtures thereof; and
- (d) from about 3-7% by weight of the composition of an alkylpolyglycoside,
- 25 the total concentration of surfactants in the composition being from about 30-35% by weight of the composition.

19. A composition according to claim 17, further comprising a cationic ammonium compound.

30

20. A composition according to claim 18, further comprising a cationic ammonium compound.

21. A composition according to claim 19, where the amounts of cationic ammonium compound is from about 0.5 to 20% by weight

of the composition.

22. A composition according to claim 20, where the amounts of cationic ammonium compound is from about 0.5 to 20% by weight of the composition.

23. A composition according to Claim 1, wherein the amount of linear alcohol ethoxylate is from about 5-40% by weight based on the weight of the composition.

24. A composition according to Claim 1, wherein the amount of linear alcohol ethoxylate is from about 5-12% by weight based on the weight of the composition.

25. An aqueous liquid cleaning composition consisting essentially of, by weight of the composition, from about 4-7% of a C_{8-10} alcohol ethoxylate having an average of about 9 moles of ethylene oxide, from about 12-20% of a C_{8-10} alcohol ethoxylate having an average of about 12 moles of ethylene oxide, from about 7-13% of a fatty acid amidopropylamine oxide having an average of about 10-18 fatty acid carbon atoms, from about 1-4% of a fatty acid diethanolamide having an average of about 10-18 fatty acid carbon atoms, from about 1-4% of a fatty acid monoethanolamide having an average of about 10-18 fatty acid carbon atoms; and an antibacterial effective amount of an antibacterial quaternary ammonium compound.

26. A composition according to claim 25, wherein the antibacterial quaternary ammonium compound is an C_8-C_{18} alkyl dimethyl benzyl ammonium chloride.

27. An aqueous liquid cleaning composition, the composition being free of anionic surfactants and comprising a nonionic surfactant system consisting essentially of:

(a) from about 1-5% by weight based on the weight of the composition of a linear alcohol ethoxylate having an average carbon chain length of from about 8 to 12 carbon atoms and from about 5-12 moles of ethoxylation per mole of alcohol;

(b) from about 3-10% by weight of the composition of a betaine;

(c) from about 1-3% by weight of the composition of a mono- or dialkanolamide; and

(d) from about 1-5% by weight of the composition of an quaternary ammonium compound.

28. An aqueous liquid cleaning composition, the composition being free of anionic surfactants and comprising a nonionic surfactant system consisting essentially of:

(a) from about 0.1-10% by weight based on the weight of the composition of a linear alcohol ethoxylate having an average carbon chain length of no more than about 12 carbon atoms and from about 4-12 moles of ethoxylation per mole of alcohol;

(b) from about 0.05-25% by weight of the composition of a betaine;

(c) from about 0.1-5% by weight of the composition of a mono- or dialkanolamide;

(d) from about 0.05 to 15% by weight of a amine oxide, the combined amount of betaine and amine oxide being from about 0.1 to 30% by weight of the composition; and

(d) from about 0.1-5% by weight of the composition of an quaternary ammonium compound,

the total concentration of surfactants in the composition no more than about 25% by weight of the composition.

29. A detergent composition suitable for preparing an aqueous, liquid cleaning formulation, the detergent composition

comprising, based on the total weight of active components in the composition:

- 5 (a) from about 10-70% by weight of an alcohol ethoxylate having an average of 6-12 carbon atoms and an average of 4-15 moles of ethylene oxide;
- (b) from about 15-60% by weight of an amine oxide, a betaine or mixture thereof;
- (c) from about 5-20% by weight of an alkanolamide having an average of 10-16 carbon atoms.

10

30. A detergent composition according to claim 29, further comprising up to about 50% by weight of a cationic surfactant.

31. An aqueous liquid composition comprising from about 3-15 10% by weight of the detergent composition of claim 29.

32. A detergent composition according to claim 29, where component (b) is an amine oxide which is an alkyldimethyl amine oxide having an average of from 10-18 carbon atoms in the alkyl
20 portion.

33. A detergent composition according to claim 29, where component (b) is an amine oxide which is an alkyl amido propyl dimethyl amine oxide having an average of from 12-18 carbon
25 atoms in the alkyl portion.

34. A detergent composition according to claim 29, where the component (b) is a betaine which is alkyl amido propyl betaine having an average of 10-18 carbon atoms in the alkyl
30 portion.

35. A detergent composition according to claim 30, wherein the cationic surfactant is a biocidal quaternary ammonium compound.

36. A detergent composition according to claim 29, further comprising an alkyl trimethyl ammonium conditioning compound.

5 37. A detergent composition according to claim 29, wherein component (b) is a combination of an amine oxide and an alkyl polyglucoside, or a betaine and an alkyl polyglucoside.

10 38. A detergent composition suitable for preparing an aqueous, liquid cleaning formulation, the detergent composition comprising, based on the total weight of active components in the composition:

- 15 a) from about 25-60% by weight an alcohol ethoxylate having an average of 6-12 carbon atoms and containing an average of from about 4-15 moles of ethylene oxide where the concentration of alcohol ethoxylate in the composition is of the total surfactant present by weight;
- 20 b) from about 15-55% by weight of an amine oxide, a betaine or mixture thereof; and
- c) 5-20% by weight of an alkanolamide having an average of from 10-16 carbon atoms; and
- d) up to about 20% by weight of a cationic surfactant.

25 39. A detergent composition according to claim 38, wherein the amount of component (b) is from about 20-35% by weight.

30 40. A detergent composition according to claim 39, wherein the cationic surfactant is from about 3-10% by weight.

 42. A detergent composition according to claim 39, wherein the alcohol ethoxylate has an average of from 8-10 carbon atoms and contains an average of 5-12 moles of ethylene oxide.

35

43. A detergent composition according to claim 39, wherein the amount of the alkanolamide in the composition is from about 9-16% by weight.

5 44. A detergent composition according to claim 39, wherein the amount of the alcohol ethoxylate is from about 25-60% by weight.

10 45. A detergent composition according to claim 39, wherein the amount of the alcohol ethoxylate is from about 40-60% by weight.

15 46. An aqueous cleaning formulation comprising from about 5% to 60% by weight of the formulation of a detergent composition,

the detergent composition comprising, based on the total weight of active components in the detergent composition:

- 20 (a) from about 25-60% by weight of an alcohol ethoxylate having an average of 8-12 carbon atoms and containing an average of 5-12 moles of ethylene oxide;
- (b) from about 15-55% by weight of an alkyl dimethyl amine oxide, an alkyl amido propyl dimethyl amine oxide, an
- 25 alkyl amido propyl betaine, or a mixture thereof;
- (c) from about 9-16% by weight of an alkanolamide having an average of 10-16 carbon atoms; and
- (d) from about 3-10% by weight of biocidal cationic surfactant.

30 47. A formulation according to claim 46, wherein the amine oxides have an average of 10-18 carbon atoms in the alkyl group.

48. A formulation according to claim 47, wherein the

35 component (b) is from about 20-35% by weight of the detergent composition.

49. A formulation according to claim 47, wherein the component (b) is an alkyl amido propyl betaine having an average of from 12-18 carbon atoms in the alkyl portion.

5

50. A composition according to claim 25, wherein the antibacterial quaternary ammonium compound is a di(C₆-C₁₄)alkyl dimethyl ammonium chloride.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 98/21788

A. CLASSIFICATION OF SUBJECT MATTER

IPC6: C11D 1/825, C11D 1/94, C11D 1/835, A61K 7/50
According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC6: C11D, A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPODOC, WPI

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	WO 9739094 A1 (STEPAN COMPANY), 23 October 1997 (23.10.97)	1-28
	--	
X	EP 0651048 A2 (EASTMAN KODAK COMPANY), 3 May 1995 (03.05.95)	1-6,9-16,23, 27,29-32, 34-35,38-39, 42-50
	--	
X	File WPI, Derwent accession no. 91-284833, Lion Corp: "Liq. detergent compsn. having good rinsability - contg. at least one non-ionic surfactant and one of amine oxide, sugar-ester type surfactant or fatty acid alkanol amide(s);" & JP,A,3188195, 910816, DW9139	1,24-9,17,29
	--	

☒ Further documents are listed in the continuation of Box C. ☒ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

10 February 1999

Date of mailing of the international search report

22. 03. 1999

Name and mailing address of the ISA/
European Patent Office, P.B. 5818 Patentlaan 2
NL-2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 cpo nl.
Fax: (+31-70) 340-3016

Authorized officer

DAGMAR JÄRVMAN

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 98/21788

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 9605807 A1 (KAO CORPORATION), 29 February 1996 (29.02.96) --	1-9,29,32, 34-36,38, 42-50
X	EP 0638637 A2 (THE PROCTER & GAMBLE COMPANY), 15 February 1995 (15.02.95) --	1-2,7,11
A	EP 0698660 A2 (KAO CORPORATION), 28 February 1996 (28.02.96) -- -----	1-4,9,10,17, 30,35,50

INTERNATIONAL SEARCH REPORT
Information on patent family members

21/12/98

International application No.
PCT/US 98/21788

Patent document cited in search report			Publication date	Patent family member(s)		Publication date
WO	9739094	A1	23/10/97	AU	2459997 A	07/11/97
EP	0651048	A2	03/05/95	CA	2132274 A	02/05/95
WO	9605807	A1	29/02/96	EP	0777466 A	11/06/97
				JP	8060188 A	05/03/96
EP	0638637	A2	15/02/95	CA	2126898 A,C	23/01/95
				US	5399280 A	21/03/95
				US	5607913 A	04/03/97
EP	0698660	A2	28/02/96	JP	7233394 A	05/09/95
				SG	32372 A	13/08/96
				US	5756443 A	26/05/98